

DIRECTORATE OF EDUCATION, GNCT OF DELHI

SUGGESTIVE ANSWERS: Term-II (2021-22)

CLASS-XI

SUBJECT: CHEMISTRY (043)

TIME: 2 Hrs.

MM: 35

1. i) a) H-bonding b) London force or Dispersion force (1/2x2)
 ii) $V_{-273} = V_0 + \frac{-273 \times V_0}{273} = 0$ (1)
2. i) Definition (1)
 ii) -ve (1)
3. i) O_2^- ion have one unpaired e^- in π_{2p}^x orbital (1)
 ii) because of its high reactivity towards air and water. (1)
4. a) $n_{CH_4} = \frac{3.2}{16} = 0.2$ $n_{CO_2} = \frac{4.4}{44} = 0.1$ (1)

Total moles = 0.2 + 0.1 = 0.3 moles

$$P = \frac{nRT}{V} = \frac{0.3 \times 0.0831 \times 300}{9} = 0.831 \text{ bar} \quad (1)$$

b) $z=1$ (1)

Or




a) $PV = nRT$ (1)

$$P = \frac{nRT}{V} = \frac{\text{mass} \times RT}{V \times M} = \frac{dRT}{M}$$



$$d = \frac{MP}{RT} \Rightarrow d \propto M$$
 (1)

(1/2 x 2=1)

b) H, He





Or



6. i) Due to inert pair effect
ii) Al become passive with conc. HNO_3 due to formation of protective layer. (1x3=3)
iii) Due to large size p- orbitals which do not cause effective overlapping.

7. i) Any two differences. (1x2=2)
ii) due to non-availability of d- orbitals can't extend its covalency beyond four. (1)


8. i) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2\text{Br}$ ii)  iii)  (1x3=3)

9. i) Any four anomalous properties
ii) $\text{Na} + (x+y)\text{NH}_3 \longrightarrow \text{Na}^+(\text{NH}_3)_x + \text{e}^-(\text{NH}_3)_y$


Or

- i) Due to small size of Be^{+2} its lattice energy with large size SO_4^{2-} in BeSO_4 is not very high and is soluble in water. Ba^{+2} being large sized, have high lattice energy large sized SO_4^{2-} in BaSO_4 and is insoluble in water. (2)
ii) $\text{Rb} < \text{K} < \text{Na} < \text{Li}$ (1)


10. a) 3- Ethyl-2- methylpentane (1)



or any another example



11. a) Target equation



On(i) x 6 + (ii) x 3 – (III)

We get $6 \times (-393.3) + 3 \times (-286.6) - (-3267.7) = 48.1 \text{ KJ}$


b- Definition

Or

- a) For spontaneous process $\Delta G < 0$ (2)
- $\Rightarrow \Delta H - T\Delta S < 0$
- $\Rightarrow 400 - T \times 0.2 < 0$
- $\Rightarrow 400 < T \times 0.2$
- $\Rightarrow T > \frac{400}{0.2} = 2000 \text{ k}$

b) 1) $\Delta n_2 = 0$ 2) $\Delta U = 0$


12 i) H^+ produced by acid combines with CH_3COO^- in solution.



ii) Definition

$$K_w = 1 \times 10^{-14} \text{ at } 298 \text{ k}$$

iii) By mixing equal moles of weak base and its salt with strong acid




$$[OH^-] = 10^{-3} \text{ M}$$

$$[H^+] = \frac{10^{-14}}{10^{-3}} = 10^{-11} \text{ M}$$

$$pH = -\log [H^+] = -\log 10^{-11} = 11$$

Or



$$[H^+] = C\alpha = C \sqrt{\frac{K_a}{C}}$$

$$[H^+] = \sqrt{K_a \cdot C}$$

Putting the values we get

$$[H^+] = 1 \times 10^{-3} \text{ M}$$

$$pH = -\log [H^+] = -\log 10^{-3}$$

$$pH = 3$$